

Optimising the use of satellite data in NWP through the inclusion of error correlation structures

L.M. Stewart, University of Reading

Co-authors: S.L. Dance and N.K. Nichols (UoR);
J.Cameron, S.English and J.Eyre (Met Office UK)

Funding: NERC and Met Office UK

Adjoint Model Applications In Dynamic Meteorology
Workshop 20 May 2009

Talk structure


- ★ Introduction and theory
 - Motivation
 - Variational data assimilation
 - Observation error covariance matrices
- ★ Quantifying observation error correlations
 - Desroziers' method of statistical approximation
 - Application to IASI data
 - Results
- ★ Modelling observation error correlation structure
 - Approximate structures for R

What are observation error correlations?

Every observation y of a atmospheric variable x has an associated error ϵ : $y = Hx + \epsilon$

→ observation error correlations are present when components of the error vector ϵ are related

→ measurement errors are attributed to 3 sources: instrument noise, forward model error and representativity error



Error sources

- **Instrument noise**
 - temperature converted $n\epsilon\delta t$ value
 - regular calibrations ensure noise is uncorrelated between channels
- **Forward model error**
 - errors in discretisation of radiative transfer equation
 - errors in mis-representation of gaseous contributors
 - errors from undetected cloud
- **Representativity error**
 - contrasting model and observation resolutions
 - observations resolve spatial scales or features that the model cannot
 - contributes to cross channel observation error correlations

Why are correlations important?

Problems

- ve magnitude and behaviour relatively unknown
- ve reduce weighting of observations in analysis
- ve for an observation vector of size 10^6 , difficult to store and invert observation error matrix if correlations are included

Benefits

- +ve increase accuracy of gradients of the observed field represented in the analysis
- +ve works with the prior error covariance to specify how observation features should be smoothed
- +ve more information available from observations

Variational data assimilation

Assimilation objective

Model forecast + Observation data \rightarrow State of atmosphere

Assimilation method Minimise a cost function which measures distance of a solution state x from the observations $y^o \in \mathbb{R}^m$ and the background field $x^b \in \mathbb{R}^n$

Cost Function

$$J(x) = \frac{1}{2}(x - x^b)^T \mathbf{B}^{-1}(x - x^b) + \frac{1}{2}(y^o - H(x))^T \mathbf{R}^{-1}(y^o - H(x))$$

where \mathbf{B} and \mathbf{R} are the background and observation error covariance matrices respectively

An error covariance matrix structure

The observation error covariance matrix takes the form:

$$\mathbf{R} = \mathbf{D}^{1/2} \mathbf{C} \mathbf{D}^{1/2}$$

where \mathbf{C} is the error correlation matrix

$$\mathbf{C} = \begin{pmatrix} 1 & \rho_{12} & \dots & \rho_{1m} \\ \rho_{12} & 1 & \dots & \rho_{2m} \\ \vdots & \ddots & \ddots & \vdots \\ \rho_{1m} & \rho_{2m} & \dots & 1 \end{pmatrix}$$

and \mathbf{D} is the error variance matrix

$$\mathbf{D} = \begin{pmatrix} \sigma^2_1 & 0 & \dots & 0 \\ 0 & \sigma^2_2 & \dots & 0 \\ \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \dots & \sigma^2_m \end{pmatrix}$$

Quantifying cross-channel correlations: a study

Objective

Generate the true observation error correlation structure for a sample set of remotely sensed data typical of NWP

Data type

- IASI (infrared atmospheric sounding interferometer) observations
- measurements of the infrared radiation emitted by the earth's surface and atmosphere at different wavelengths

Method

We use a post analysis diagnostic derived from variational data assimilation theory [Desroziers, 2005]

Desroziers' method of statistical approximation

Recall the background state, x_b , and observation vector, y , are approximations to the true state of the atmosphere, x_t ,

$$\begin{aligned}y &= Hx_t + \epsilon^o \\x_t &= x_b + \epsilon^b\end{aligned}$$

where ϵ^o and ϵ^b are the observation and background errors respectively.

The best linear unbiased estimate of the true state, x_a , is given by

$$\begin{aligned}x_a &= x_b + \mathbf{K}(y - Hx_b) = x_b + \mathbf{K}d_b^o \\ \mathbf{K} &= \mathbf{B}\mathbf{H}^T(\mathbf{H}\mathbf{B}\mathbf{H}^T + \mathbf{R})^{-1}\end{aligned}$$

Desroziers' method of statistical approximation

Innovation vector

$$\begin{aligned}d_b^o &= y - Hx_b = Hx_t + \epsilon^o - Hx_b \\ &\approx \epsilon^o + \mathbf{H}\epsilon^b\end{aligned}$$

Analysis innovation vector

$$\begin{aligned}d_a^o &= y - Hx_a = y - H(x_b + Kd_b^o) \\ &\approx (\mathbf{I} - \mathbf{HK})d_b^o \\ &\approx \mathbf{R}(\mathbf{HBH}^T + \mathbf{R})^{-1}d_b^o\end{aligned}$$

Desroziers' method of statistical approximation

Taking the expectation of the cross product of d_a^o and d_b^o , and assuming

$$\mathbb{E}[\epsilon^o(\epsilon^b)^T] = \mathbb{E}[\epsilon^b(\epsilon^o)^T] = 0,$$

we find a statistical approximation for the observation error covariances

$$\begin{aligned} \mathbb{E} [d_a^o(d_b^o)^T] &\approx \mathbb{E} [\mathbf{R}(\mathbf{H}\mathbf{B}\mathbf{H}^T + \mathbf{R})^{-1}d_b^o(d_b^o)^T] \\ &\approx \mathbf{R}(\mathbf{H}\mathbf{B}\mathbf{H}^T + \mathbf{R})^{-1}\mathbb{E} [(\epsilon^o + H\epsilon^b)(\epsilon^o + H\epsilon^b)^T] \\ &\approx \mathbf{R}(\mathbf{H}\mathbf{B}\mathbf{H}^T + \mathbf{R})^{-1}(\mathbf{R} + \mathbf{H}\mathbf{B}\mathbf{H}^T) \\ &\approx \mathbf{R} \end{aligned}$$

Application to IASI data




Figure: Assimilation process

Observation error correlation matrix






Figure: Error correlation matrix for 139 channels used in Var

Observation error correlation matrix



(a)



(b)

Figure: Error correlation matrix for (a) temperature sounding channels; (b) water vapour channels

Operational and diagnosed error variances




Figure: Operational error variances (black line), diagnosed error variances (red line), and first off-diagonal error covariance (green line)

Diagnosed error variances: comparison with Hollingsworth-Lonnberg (H-L) method




Figure: Diagnosed error variances (red line), H-L diagnosed error variances for 84.8km (blue and black line) and 61.9km (green line) separation. Plot provided by James Cameron, UK Met Office.

Quantifying cross-channel correlations: a summary

- ★ Strong off-diagonal correlations are present between channels with similar spectral properties
- ★ Channels highly sensitive to water vapour have large observation error variances and covariances
- ★ The observation error variance is being overestimated in current assimilation algorithms
- ★ Diagnosed error variances are comparable with those using the H-L diagnostic
- ★ Non-symmetric matrices! → future work

Modelling error correlation structure

What next?

Investigate how to approximate the true error correlation structure within operational assimilation methods...

Current approaches

- a diagonal matrix approximation
- diagonal variance inflation

Alternative approaches

- a Markov error covariance approximation
- a truncated eigendecomposition approximation [Fisher, 2005]
- a Toeplitz to circulant matrix approximation [Healy, 2005]

A desirable error covariance matrix

Main issue in observation error correlation modelling

- ★ need to calculate matrix-vector product $\mathbf{R}^{-1}(y^o - Hx)$ every time we calculate cost function J
- ★ relatively easy if $\mathbf{R} = \mathbf{D} \equiv m$ scalar multiplications
- ★ **BUT** $y \in \mathbb{R}^{10^6}$ and so $\mathbf{R} \in \mathbb{R}^{10^6 \times 10^6}$ which, if dense, is impossible to store and invert

The perfect partner: what do we want from $\mathbf{R} \neq \mathbf{D}$?

- ♥ structure resulting in an \mathbf{R}^{-1} suitable for storage / can be used cheaply in a matrix-vector product
- ♥ representative of the true error correlation structure
- ♥ greater access to information from the observations and improved analysis accuracy

A Markov error covariance approximation

Consider a Markov covariance matrix of the form

$$R_{ij} = \sigma^2 \rho^{|i-j|}, \quad \rho = \exp\left(-\frac{\delta z}{h}\right)$$

where σ^2 is the error variance, δz is the level spacing, and h is the length scale

This is equivalent to a correlation matrix of the form

$$C = \begin{pmatrix} 1 & \rho & \rho^2 & \dots & \rho^n \\ \rho & 1 & \rho & \dots & \rho^{n-1} \\ \rho^2 & \rho & 1 & \dots & \rho^{n-2} \\ \vdots & \vdots & \vdots & & \vdots \\ \rho^n & \dots & \rho^2 & \rho & 1 \end{pmatrix}$$

A Markov error covariance approximation

The benefit of this choice is that C has a tri-diagonal inverse

$$C^{-1} = \frac{1}{1 - \rho^2} \begin{pmatrix} 1 & -\rho & 0 & \dots & 0 \\ -\rho & 1 + \rho^2 & -\rho & \dots & 0 \\ 0 & -\rho & 1 + \rho^2 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \dots & -\rho & 1 + \rho^2 & -\rho \\ 0 & \dots & 0 & -\rho & 1 \end{pmatrix}$$

and therefore as does R : $R^{-1} = \frac{1}{\sigma^2} I \times C^{-1}$

No need to store and invert R !

An eigendecomposition approximation

Describe C by a truncated eigendecomposition using its leading eigenpairs

$$\tilde{R} = D^{1/2} \tilde{C} D^{1/2} = D^{1/2} \left(\alpha I + \sum_{k=1}^K (\lambda_k - \alpha) \mathbf{v}_k \mathbf{v}_k^T \right) D^{1/2}$$

where $(\lambda_k, \mathbf{v}_k)$ is an eigenvalue, eigenvector pair of C , K is the number of eigenpairs used, and α is chosen such that $\text{trace}(\tilde{R}) = \text{trace}(D)$ [Fisher, 2005]

This matrix also has an easily attainable inverse

$$\tilde{R}^{-1} = D^{-1/2} \left(\alpha^{-1} I + \sum_{k=1}^K (\lambda_k^{-1} - \alpha^{-1}) \mathbf{v}_k \mathbf{v}_k^T \right) D^{-1/2}$$

No need to store and invert R !

Summary

- ★ Observation error correlations are often created because of contrasting model and observation resolutions
- ★ Including observation error correlation structure can increase analysis accuracy and information content
- ★ In IASI data, observation error correlations are strongest between channels with similar spectral properties
- ★ In IASI data, the largest observation error covariances are between channels highly sensitive to water vapour
- ★ In order to include observation error correlation structure in data assimilation algorithms, the **R** matrix must be suitably structured

References

- G. Desroziers and L. Berre and B. Chapnik and P. Poli. *Diagnosis of observation, background and analysis-error statistics in observation space*. Q.J.R.Meteorol.Soc., 131, 2005.
- M. Fisher. *Accounting for Correlated Observation Error in the ECMWF Analysis*. ECMWF Technical Memoranda, MF/05106, 2005.
- S.B. Healy and A.A. White. *Use of discrete Fourier transforms in the 1D-Var retrieval problem*. Q.J.R.Meteorol.Soc., 131, 2005.